

IN THE CLAIMS


Please amend the claims as follows:

Claims 1-6 (Cancelled)

Claim 7 (Currently Amended): A method for producing an allyl compound comprising:

reacting a first allyl compound of formula (a) with an oxygen nucleophilic agent in the presence of a catalyst containing a monodentate phosphite compound and at least one transition metal compound containing a transition metal selected from the group consisting of transition metals belonging to Group 8 to Group 10 of the Periodic Table, under conditions suitable for producing a second allyl compound;


wherein the monodentate phosphite compound has the following formula (1):



wherein R^1 , R^2 and R^3 are ~~respectively~~ independently an alkyl group which may have a substituent, a carbon chain having of R^1 , R^2 and R^3 may have at least one carbon-carbon double bond or triple bond, or at least two of R^1 , R^2 and R^3 bond to each other to form a cyclic structure containing oxygen and at least two optional groups of R^1 , R^2 and R^3 may bond to each other to form at least one cyclic structure;

wherein the first allyl compound has the following formula (a):

formula (a):



wherein R^a to R^e are respectively independently a hydrogen atom, a halogen atom, a hydroxyl group, an amino group, a formyl group, an alkyl group, an aryl group (including a

heterocyclic compound forming an aromatic 6π electron cloud on the upper and lower sides of the ring, hereinafter the same), an alkoxy group, an aryloxy group, an alkylthio group, an arylthio group, an amide group, an acyl group or an acyloxy group; among these groups, the amino group, the alkyl group, the aryl group, the alkoxy group, the aryloxy group, the alkylthio group, the arylthio group, the acyl group or the acyloxy group may have a substituent; and when any of R^a to R^e has a carbon chain, the carbon chain may have at least one carbon-carbon double bond or triple bond;

X is a halogen atom, a hydroxyl group, a nitro group, an amino group, a sulfonyl group, a sulfonate group, an acyloxy group, a carbonate group, a carbamate group, a phosphate group, an alkoxy group or an aryloxy group; among these groups, the amino group, the sulfonyl group, the sulfonate group, the acyloxy group, the carbonate group, the carbamate group, the phosphate group, the alkoxy group and the aryloxy group may have a substituent; when X has a carbon chain, the carbon chain may have at least one carbon-carbon double bond or triple bond; and at least two optional groups among R^a to R^e and X may bond to each other to form at least one cyclic structure; and

wherein the second allyl compound has a compositional structure different than that of the first allyl compound,

wherein the oxygen nucleophilic agent and X are not a monodentate phosphite ~~is not the same as the monodentate phosphite compound and is not the same as the X of the first allyl compound, X^- , or the proton adduct of X.~~

Claim 8 (Previously Presented): The method of Claim 7, wherein the first allyl compound has a molecular weight of 1,500 Da or less or contains at most 100 carbon atoms.

Claim 9 (Previously Presented): The method of Claim 7, wherein R^a to R^e are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, aryl, and substituted aryl.

Claim 10 (Previously Presented): The method of Claim 7, wherein X is a halogen atom.

Claim 11 (Previously Presented): The method of Claim 7, wherein X is a hydroxy group, an alkoxyl group or an aryloxy group.

Claim 12 (Previously Presented): The method of Claim 7, wherein X is a nitro group or an amino group.

Claim 13 (Previously Presented): The method of Claim 7, wherein X is a sulfonyl group or a sulfonate group.

Claim 14 (Previously Presented): The method of Claim 7, wherein X is an acyloxy group, a carbonate group, a carbamate group, or a phosphate group.

Claim 15 (Previously Presented): The method of Claim 7, wherein the oxygen nucleophilic agent is an alcohol, a deprotonated alcohol, a hydroxaryl, a deprotonated hydroxaryl, an aromatic carboxylic acid, or a deprotonated aromatic carboxylic acid.

Claim 16 (Previously Presented): The method of Claim 7, wherein the catalyst contains a monodentate phosphite compound of formula (I), wherein at least one of R^1 , R^2

and R³ of the monodentate coordinated phosphite compound of formula (1) is a branched chain-like alkyl group.

Claim 17 (Cancelled)


Claim 18 (Previously Presented): The method of Claim 7, wherein the catalyst contains a monodentate phosphite compound of formula (I), wherein R¹, R² and R³ are each independently an alkyl group which may have a substituent.

Claim 19 (Currently Amended): The method of Claim 7, wherein the catalyst contains a monodentate phosphite compound of formula (I), wherein at least one of R¹, R² and R³ ~~is an alkyl group which~~ has a carbon-carbon double bond or triple bond.


Claim 20 (Currently Amended): The method of Claim 7, wherein the catalyst contains a monodentate phosphite compound of formula (I), wherein at least two of R¹, R² and R³ bond to each other to form at least one cyclic structure containing oxygen.

Claim 21 (Previously Presented): The method of Claim 7, wherein said monodentate compound is selected from the group consisting of compounds of formulas L-1, L-2, L-3, L-4, L-5, L-6, L-7, L-8, L-9, and L-10;


wherein formulas L-1, L-2, L-3, L-4, L-5, L-6, L-7, L-8, L-9, and L-10 are:




(L-1)




(L-2)




(L-3)




(L-4)




(L-5)




(L-6)




(L-7)



(L-8)



(L-9)



(L-10)

Claim 22 (Previously Presented): The method of Claim 7, wherein the transition metal compound is at least one compound selected from the group consisting of a ruthenium compound, a rhodium compound, an iridium compound, a nickel compound, a palladium compound and a platinum compound.

Claim 23 (Previously Presented): The method of Claim 7, wherein the transition metal compound is a palladium compound.

Claim 24 (Previously Presented): The method of Claim 7, wherein the oxygen nucleophilic agent is a compound different from a substituent X and its proton adduct X-H is eliminated from the allyl starting material compound by reaction, and is a compound containing an oxygen atom expressed by AO-H or its deprotonated form of AO⁻, in which A is a hydrogen atom or an organic group having a carbon atom, a nitrogen atom, a phosphorus atom or a sulfur atom bonded to an oxygen atom.

Claim 25 (Previously Presented): The method of Claim 7, wherein a phosphonium compound is present during the reaction.

Claim 26 (Previously Presented): The method of Claim 7, wherein an ammonium compound is present during the reaction.

Claim 27 (New): The method of Claim 7, wherein R₁, R₂ and R₃ are linear alkyl, alkenyl or alkynyl groups.

Claim 28 (New): The method of Claim 7, wherein R_1 , R_2 and R_3 are linear alkyl groups.

Claim 29 (New): The method of Claim 7, wherein R_1 , R_2 and R_3 are linear C_1 - C_{10} alkyl groups.